

STI WARNING SIGNS

RECCOMENDATIONS: It is recommended that you see a healthcare provider for any of the following symptoms that might indicate an STD:

- Anal pain or itching (butthole hurts or itches)
 Itching in your pubic area (privates), more than 2 weeks
- Lice or lice eggs seen in your pubic area (privates)
- ↓ Sore or bump on genitals (privates) in past 12 months
- ↓ Widespread rash, both sides body, past 12 months
- ↓ Female with abnormal vaginal discharge (odd smell or color), over 5 days
- Female with bleeding after sex
- ✤ Female with daily lower abdominal pain, not during period
- **4** Male with discharge from penis or burning during urination
- 4 Male with severe testicular pain.

IF YOU DO NOT START YOUR PERIOD, PLEASE NOTE THE FOLLOWING PREGNANCY WARNING SIGNS

Call your doctor or go to a local urgent care or ER for the following:

- 1. Vaginal Bleeding: Some spotting is normal, but bleeding could be a sign of miscarriage or ectopic pregnancy. Heavy bleeding, cramping or abdominal pain is not the norm, and may signal a miscarriage or ectopic pregnancy
- 2. Excessive Nausea and Vomiting: You are unable to keep down food or fluids.
- 3. High Fever: A fever greater than 101 degrees during pregnancy may be serious.
- 4. Pain and Burning during Urination: These can be signs of bladder or urinary tract infections, which need prompt attention.